

AD-A141 652

ON THE RANGE OF A REGENERATIVE SEQUENCE(U) WISCONSIN  
UNIV-MADISON MATHEMATICS RESEARCH CENTER P W GLYNN  
MAR 84 MRC-TSR-2661 DAAG29-80-C-0041


1/1

UNCLASSIFIED

F/G 12/1

NL

END  
DATE  
FILMED  
7-83  
DTIC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS 1963 A

AD-A141 652

MRC Technical Summary Report # 2661

ON THE RANGE OF A REGENERATIVE SEQUENCE

Peter W. Glynn

Mathematics Research Center  
University of Wisconsin—Madison  
610 Walnut Street  
Madison, Wisconsin 53705

March 1984

(Received November 30, 1983)

DTIC FILE COPY

Approved for public release  
Distribution unlimited

Sponsored by

U. S. Army Research Office  
P. O. Box 12211  
Research Triangle Park  
North Carolina 27709

DTIC  
ELECTE  
MAY 31 1984

84 05 31 073

UNIVERSITY OF WISCONSIN - MADISON  
MATHEMATICS RESEARCH CENTER

ON THE RANGE OF A REGENERATIVE SEQUENCE

Peter W. Glynn

Technical Summary Report #2661

March 1984

ABSTRACT

For a given countable partition of the range of a regenerative sequence  $\{X_n : n \geq 0\}$ , let  $R_n$  be the number of distinct sets in the partition visited by  $X$  up to time  $n$ . We study convergence issues associated with the range sequence  $\{R_n : n \geq 0\}$ . As an application, we generalize a theorem of Chosid and Isaac to Harris recurrent Markov chains.

AMS (MOS) Subject Classifications: 60K05, 60J27, 60F15

Key Words: regenerative process, range, subadditive process, Markov chain.

Work Unit Number 5 - Optimization and Large Scale Systems

# SIGNIFICANCE AND EXPLANATION

Consider a system which evolves randomly in time; the trajectory of such a system traces a path through space. If one partitions space into a disjoint collection of subsets, one can study the number of subsets visited by the trajectory up to a certain instant. In this paper, we show that, under certain conditions, the number of subsets never grows linearly in time, regardless of the partition used. On the other hand, the precise order of growth (which can be arbitrarily close to linear order) does depend on the way in which space is partitioned. These results are obtained for regenerative random processes. Such processes describe systems which, when viewed on a certain random time scale, evolve in an independent and identically distributed fashion. Virtually any ergodic discrete-time Markov chain has this property.

Accession For	
GRA&I	<input checked="" type="checkbox"/>
TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
For	
Distribution/	
Availability Codes	
Distribution/	
Dist. Special	
A-1	



The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

# ON THE RANGE OF A REGENERATIVE SEQUENCE

Peter W. Glynn

## 1. Introduction

Let  $\{X_n : n \geq 0\}$  be a stochastic sequence taking values in a measurable space  $(E, \mathcal{E})$ . For a given family  $\{A_n : n \geq 0\}$  of  $\mathcal{E}$ -measurable sets partitioning  $E$ , set

$$Y_n = \sum_{k=0}^{\infty} k I(X_n \in A_k) ,$$

where  $I(\Lambda)$  is 1 or 0 depending on whether or not  $\omega \in \Lambda$ . Let  $\phi(1,1) = 1$  and set

$$\phi(1,j) = I(Y_1 \neq Y_j, Y_{1+1} \neq Y_j, \dots, Y_{j-1} \neq Y_j)$$

$$(1.1) \quad R(1,j) = \sum_{k=1}^j \phi(1,k)$$

for  $1 < j$ . The process  $\{R(0,n) : n \geq 0\}$ , which is called the range sequence associated with  $\{X_n : n \geq 0\}$ , counts the number of distinct sets  $A_k$  visited by  $X$  up to time  $n$ . The range process  $\{R(0,n) : n \geq 0\}$  has been extensively studied, in the case that  $\{X_n : n \geq 0\}$  is a random walk with stationary increments; see, for example, Dvoretzky and Erdos (1951), p. 35-40 of Spitzer (1976), and Jain and Pruitt (1972). Chosid and Isaac (1978, 1980) have considered the problem when  $\{X_n : n \geq 0\}$  is a recurrent countable state Markov chain.

In this paper, we wish to study the range process in the case that  $\{X_n : n \geq 0\}$  is a delayed regenerative sequence, with associated regeneration times  $T_1, T_2, \dots$  (see p. 298-302 of Çinlar (1975) for the definition). Our starting point is a simple decomposition formula for the range. In Section 2, we use this formula to improve the main result of Chosid and Isaac. Section 3 discusses the application of the results to Harris recurrent Markov chains, and considers a generalization to the case where  $\{A_n : n \geq 0\}$  is not a partition of  $E$ .

## 2. Convergence Results for the Range

Let  $R_{ik} = R(T_i, T_k - 1)$  for  $i < k$ ,  $W_i = R_{1,i+1} - R_{1i}$ , and set  $S(n) = \inf\{j > T_1 : Y_j = n\}$ . Then,

$$\begin{aligned}
 W_i &= \sum_{k=T_i}^{T_{i+1}-1} \phi(T_1, k) \\
 &= \sum_{k=T_i}^{T_{i+1}-1} \phi(T_1, k) \cdot \sum_{n=0}^{\infty} I(Y_k = n) \\
 (2.1) \quad &= \sum_{n=0}^{\infty} \sum_{k=T_i}^{T_{i+1}-1} I(Y_{T_1} = n, \dots, Y_{k-1} = n, Y_k = n) \\
 &= \sum_{n=0}^{\infty} I(T_i < S(n) < T_{i+1}) .
 \end{aligned}$$

The decomposition formula (2.1) plays an important role in our development.

(2.2) Lemma. i.) Set  $\tau_i = T_{i+1} - T_i$ . Then  $E(W_i/\tau_i) \rightarrow 0$  as  $i \rightarrow \infty$ .

ii.) If  $EW_1^m < \infty$ , then  $EW_i^m \rightarrow 0$  as  $i \rightarrow \infty$ .

Proof. For i.), observe that

$$\begin{aligned}
 E(W_i/\tau_i) &= \sum_{n=0}^{\infty} E\{1/\tau_i ; T_i < S(n) < T_{i+1}\} \\
 (2.3) \quad &= \sum_{n=0}^{\infty} P\{S(n) > T_2\}^{i-1} E\{1/\tau_1 ; S(n) < T_2\}
 \end{aligned}$$

where the second equality follows from the regenerative property of  $X$ . Since

$$\sum_{n=0}^{\infty} E\{1/\tau_1 ; S(n) < T_2\} = E(W_1/\tau_1) < 1 ,$$

one may apply bounded convergence to (2.3) to obtain i.). For ii.), note that

$$\begin{aligned}
EW_1^m &= \sum_{n_1, \dots, n_m} P\{T_1 < S(n_1), \dots, S(n_m) < T_{i+1}\} \\
(2.4) \quad &= \sum_{n_1, \dots, n_m} P\{S(n_1) > T_2, \dots, S(n_m) > T_2\}^{i-1} \cdot P\{S(n_1) < T_2, \dots, S(n_m) < T_2\}
\end{aligned}$$

The bounded convergence theorem then applies to (2.4) provided that

$$\sum_{n_1, \dots, n_m} P\{S(n_1) < T_2, \dots, S(n_m) < T_2\} = EW_1^m < \infty. \quad ||$$

By Minkowski's inequality

$$\begin{aligned}
(E(R_{1n}/n)^m)^{1/m} &< \sum_{i=1}^{n-1} (E(W_1/n)^m)^{1/m} \\
&= \frac{1}{n} \sum_{i=1}^{n-1} (EW_1^m)^{1/m}
\end{aligned}$$

so the following corollary is immediate.

(2.5) Corollary. If  $EW_1^m < \infty$ , then  $E(R_{1n}/n)^m \rightarrow 0$  as  $n \rightarrow \infty$ .

The following lemma shows that the process  $\{R_{ik} : 0 \leq i < k\}$  ( $T_0 = 0$ ) obeys a sub-additive inequality.

(2.6) Lemma. If  $0 \leq i < j < k$ , then  $R_{ik} \leq R_{ij} + R_{jk}$ .

Proof. If  $0 \leq i < j < k$ , then (1.1) implies that

$$\begin{aligned}
R_{ik} &= R_{ij} + \sum_{\ell=T_j}^{T_k-1} \phi(T_i, \ell) \\
&= R_{ij} + \sum_{\ell=T_j}^{T_k-1} I(Y_{T_i} \neq Y_\ell, \dots, Y_{\ell-1} \neq Y_\ell) \\
&\leq R_{ik} + \sum_{\ell=T_j}^{T_k-1} I(Y_{T_j} \neq Y_\ell, \dots, Y_{\ell-1} \neq Y_\ell) \\
&= R_{ij} + R_{jk}. \quad ||
\end{aligned}$$

The following theorem now follows easily.

(2.7) Theorem. If  $EW_1 < \infty$ , then  $R_{0n}/n \rightarrow 0$  a.s.

Proof. The regenerative structure of  $X$  implies that the distribution of  $\{R_{k+1,j+1} : 1 \leq k < j\}$  is identical to that of  $\{R_{kj} : 1 \leq k < j\}$ . By virtue of Corollary 2.5 ( $m = 1$ ), the non-negativity of  $R_{kj}$ , and Lemma 2.6, this implies that  $\{R_{kj} : 1 \leq k < j\}$  satisfies the postulates of the subadditive ergodic theorem (see Kingman (1973)). Hence,  $R_{1n}/n \rightarrow 0$  as  $n \rightarrow \infty$  a.s. But by Lemma 2.6,

$$0 \leq R_{0n}/n \leq R_{01}/n + R_{1n}/n \rightarrow 0 \text{ a.s.}$$

as  $n \rightarrow \infty$ . ||

We wish to point out that the subadditive ergodic theorem has been previously used to analyze the mean range in a different setting; see Derriennic (1980).

Corollary 2.5, Lemma 2.6, and the observation that  $R(0,n) \leq R_{0,n+1}$  yield the next result.

(2.8) Corollary i.) If  $EW_1 < \infty$ , then  $R(0,n)/n \rightarrow 0$  a.s.

ii.) If  $E(R_{01} + W_1)^m < \infty$ , then  $E(R_{0n}/n)^m \rightarrow 0$  as  $n \rightarrow \infty$ .

Recall that any irreducible recurrent Markov chain  $\{X_n : n \geq 0\}$  on  $\{0, 1, \dots\}$  can be regarded as a regenerative sequence. In particular, Corollary 2.8 shows that if

$$\alpha(i) \triangleq E \left[ \sum_{j=0}^{T_1(i)-1} \phi(0,j) \mid X_0 = i \right] < \infty$$

( $T_n(i) = \inf\{m > T_{n-1}(i) : X_m = i\}$ ,  $T_0(i) = 0$ ) for some  $i$ , then  $R(0,n)/n \rightarrow 0$  a.s. This is Theorem 1 of [1]. Note that if  $\{X_n : n \geq 0\}$  is positive recurrent, then  $\alpha(i) < \infty$ , so that the mean range then automatically converges to zero a.s. Our results are somewhat stronger than those of [1] in the null recurrent case. Theorem 2.7 proves that  $R_{0n}/n \rightarrow 0$  a.s., whereas [1] only proves that  $R_{0n}/n$  is a.s. bounded under the hypothesis  $EW_1 < \infty$ . In the same spirit, Corollary 2.8 ii.) and Lemma 2.2 i.) are more complete than those of [1].

A natural question, in the context of the Markov chains discussed above, is whether finiteness of  $\alpha(i)$  is a solidarity property; in other words, if  $\alpha(i)$  is finite for one  $i$ , need  $\alpha(i)$  be finite for all  $i$ . Of course, if  $R(0,n)/n \rightarrow 0$  a.s. were to imply that  $\alpha(i) < \infty$ , we would be done. However, as pointed out in [2], a symmetric nearest neighbor random walk on the integers obeys  $R(0,n)/n \rightarrow 0$  a.s., and yet  $\alpha(i) = \infty$  for all  $i$ .

Let  $E_j(\cdot)$ ,  $P_j(\cdot)$  denote expectation and probability respectively, conditional on  $X_0 = j$ . Let

$$V(i,j) = \inf\{n \geq 1 : T_{n-1}(j) \leq T_1(i) < T_n(j)\}.$$

(2.9) Lemma. Let  $X$  be an irreducible recurrent Markov chain. Then, for any  $i$  and  $j$ ,  $\alpha(j) < \infty$  if and only if  $E_j R(0, T_{V(i,j)}(j) - 1) < \infty$ .

Proof. By Lemma 2.6,

$$\begin{aligned} R(0, T_{V(i,j)}(j) - 1) &\leq \sum_{k=1}^{V(i,j)} R(T_{k-1}, T_k - 1) \\ (2.10) \quad &= \sum_{k=1}^{\infty} R(T_{k-1}, T_k - 1) I(V(i,j) \geq k) \\ &+ \sum_{k=1}^{\infty} R(T_{k-1}, T_k - 1) I(V(i,j) = k). \end{aligned}$$

Taking expectations in (2.10) and using the regenerative property yields

$$\begin{aligned} E_j R(0, T_{V(i,j)}(j) - 1) &\leq \sum_{k=1}^{\infty} P_j\{T_1(i) \geq T_1(j)\}^{k-1} E_j\{R(0, T_1(j) - 1) : T_1(i) \geq T_1(j)\} \\ &+ \sum_{k=1}^{\infty} P_j\{T_1(i) \geq T_1(j)\}^{k-1} E_j\{R(0, T_1(j) - 1) : T_1(i) < T_1(j)\} \\ &= \alpha(j) / P_j\{T_1(i) < T_1(j)\} \end{aligned}$$

which proves one implication. For the other, note that  $T_1(j) \leq T_{V(i,j)}(j)$ . ||

(2.11) Theorem. Let  $X$  be an irreducible recurrent Markov chain. Then, if  $\alpha(i) < \infty$  for one  $i$ ,  $\alpha(i)$  is finite for all  $i$ .

Proof. By Lemma 2.9, we will be done if we prove that

$$(2.12) \quad E_j R(0, T_{V(i,j)}(j)) - 1 = E_i R(0, T_{V(j,i)}(i)) - 1.$$

By the same reasoning which produced (2.1),

$$(2.13) \quad R(0, T_{V(i,j)}(j)) - 1 = \sum_{n=0}^{\infty} I(\tilde{S}(n) < T_{V(i,j)}(j)).$$

where  $\tilde{S}(n) = \inf\{k > 0 : Y_k = n\}$ . But by the strong Markov property applied at  $T_1(i)$ ,

$$\begin{aligned} P_j\{S(n) < T_{V(i,j)}(j)\} \\ &= P_j\{S(n) < T_1(i)\} + P_j\{S(n) > T_1(i)\} P_1\{S(n) < T(j)\} \\ &= P_1\{S(n) < T_1(j)\} + P_1\{S(n) > T_1(j)\} P_j\{S(n) < T(i)\} \\ &= P_1\{S(n) < T_{V(j,i)}(i)\}. \end{aligned}$$

Substituting in (2.13) yields the equality (2.12). ||

It should be mentioned that a similar argument appears in Chung (1966), p. 84, in connection with a solidarity problem concerning moments of certain functionals.

As indicated in [1], the mean range  $R(0,n)/n$  can display a wide range of different limit behavior, in the absence of the moment condition  $EW_1 < \infty$ .

(2.14) Example. Let  $\tau_1, \tau_2, \dots$  be a sequence of positive independent identically distributed integer-valued random variables. Suppose their common distribution  $F$  is in the domain of attraction of a stable law with parameter  $0 < \alpha < 1$ . Let  $S_n = \tau_1 + \dots + \tau_n$  and set  $l(n) = \max\{k : S_k \leq n\}$ . The process  $X_n = S_{l(n)+1} - n$  is an irreducible recurrent Markov chain on  $\{0, 1, 2, \dots\}$ . Letting  $A_n = \{n\}$ , we consider the mean range on the subsequence  $\{S_n : n \geq 1\}$ . It is easily checked that

$$R(0, S_n - 1)/S_n = \max_{1 \leq k \leq n} \tau_k/S_n$$

so that

$$R(0, S_{n-1})/S_n \Rightarrow Z(\alpha)$$

as  $n \rightarrow \infty$ , where  $Z(\alpha)$  is a non-degenerate r.v. and  $\Rightarrow$  denotes weak convergence (see Feller (1971), p. 465).

So far, our study of convergence has centered on normalizing  $R(0,n)$  by  $n$ . In view of the geometric factors in (2.3) and (2.4), it is natural to investigate whether normalizing by  $n^{-\alpha}$  ( $0 < \alpha < 1$ ) is adequate for  $ER(0,n)$ .

(2.15) Example. Let  $\{X_n : n \geq 0\}$  be a sequence of non-negative independent identically distributed integer-valued random variables. Then,  $X$  is regenerative with  $T_n = n$ , and  $EW_1 < 1$ . Suppose that  $n^{-\alpha}ER(0,n) \rightarrow 0$  as  $n \rightarrow \infty$  where  $\alpha < 1$ . Then,

$$\begin{aligned} \sum_{k=1}^n E\phi(0,k)/k &= \sum_{k=1}^n (ER(0,k) - ER(0,k-1))/k \\ (2.16) \quad &= \sum_{k=1}^n ER(0,k)/k(k+1) + R(0,n)/(n+1) - R(0,0) ; \end{aligned}$$

letting  $n \rightarrow \infty$ , we see that the second term vanishes and the first is summable since we are assuming that  $ER(0,n) = O(n^\alpha)$  for  $\alpha < 1$ . We will now show that (2.16) diverges in general, proving that  $ER(0,n) = O(n^\alpha)$  is not valid without further assumptions. Using (2.1), we see that

$$\begin{aligned} \sum_{k=1}^{\infty} E\phi(0,k)/k &= \sum_{n=0}^{\infty} \sum_{k=1}^{\infty} P\{X_1 \neq n\}^{k-1} P\{X_1 = n\}/k \\ (2.17) \quad &= - \sum_{n=0}^{\infty} \log(P\{X_1 = n\})P\{X_1 = n\} \end{aligned}$$

where we interpret  $\log(0) \cdot 0 = 0$ . Choosing the mass function  $P\{X_1 = n\}$  to be  $C/n(\log n)^2$  (some constant  $C$ ) for  $n \geq 2$  causes (2.17) to diverge.

We therefore have the following conclusion. If  $\{X_n : n \geq 0\}$  is a regenerative sequence for which  $EW_1 < \infty$ , then  $ER(0,n)/n$  converges to zero regardless of the partition used. On the other hand, one needs conditions on the partition to ensure convergence of  $ER(0,n)/n^\alpha$  to zero, for  $\alpha < 1$ .

(2.18) Proposition. Suppose that  $\sum_{n=0}^{\infty} n^\gamma P\{S(n) < T_2\} < \infty$  for some  $\gamma > 1$ . Then,  $n^{-\alpha}R_{0n} \rightarrow 0$  a.s. and  $n^{-\alpha}ER_{1n} \rightarrow 0$ , as  $n \rightarrow \infty$ , where  $\alpha = 1/\gamma$ .

Proof. Let  $Z_n = \max\{k : S(k) < T_{n+1}\}$  and observe that  $R_{1n} < Z_n$ . But  $Z_n = \max\{v_1, \dots, v_n\}$ , where  $\{v_i : i \geq 1\}$  is the independent and identically distributed sequence defined by

$$(2.19) \quad v_i = \max\{k : T_i < S(k) < T_{i+1}\}.$$

Noting that

$$(2.20) \quad \begin{aligned} EV_1^Y &= \sum_{n=0}^{\infty} n^Y P\{S(n) < T_2 ; S(m) > T_2 \text{ for } m > n\} \\ &< \sum_{n=0}^{\infty} n^Y P\{S(n) < T_2\}, \end{aligned}$$

it follows that  $EV_1^Y < \infty$  under our hypotheses. Hence,  $\sum_{k=1}^n v_k^Y/n$  converges to a finite quantity, which implies that  $Z_n^Y/n \rightarrow 0$  a.s. Thus,  $n^{-\alpha}R_{0n} \rightarrow 0$  a.s.

For the convergence of  $n^{-\alpha}ER_{1n}$ , we use the finiteness of  $EV_k^Y$  to conclude that  $EZ_n^Y/n \rightarrow 0$  as  $n \rightarrow \infty$  (see p. 90 of Chung (1967)). But

$$(2.21) \quad 0 < (ER_{1n}/n^\alpha)^Y < (EZ_n^Y/n^\alpha)^Y < EZ_n^Y/n,$$

where the last inequality is a statement of the fact that  $(E|X|^r)^{1/r}$  is a non-decreasing function of  $r$  for any random variable  $X$ . (Feller (1971), p. 155.) Relation (2.21) implies our result. ||

Recalling that  $R(0,n) \leq R_{01} + R_{1n}$ , we obtain the following corollary.

(2.22) Corollary. Assume that the conditions of Proposition 2.18 are in force. If  $ER_{01} < \infty$ , then  $n^{-\alpha}ER(0,n) \rightarrow 0$  as  $n \rightarrow \infty$ , for  $\alpha > 1/\gamma$ .

To conclude this section, recall that if  $\{X_n : n \geq 0\}$  is a delayed regenerative sequence for which  $ET_1 < \infty$ , then

$$\frac{1}{n} \sum_{k=1}^n P\{X_k \in B\} \rightarrow \pi(B) \triangleq \frac{1}{ET_1} P\{X_{T_1+k} \in B; \tau_1 > k\}$$

as  $n \rightarrow \infty$  (see [4], p. 299); the probability  $\pi$  is called the ergodic measure for the sequence. Note that

$$\begin{aligned} P\{S(n) < T_2\} &< \sum_{k=0}^{\infty} P\{X_{T_1+k} \in A_n; \tau_1 > k\} \\ &= \pi(A_n)ET_1. \end{aligned}$$

Hence, if  $Et_1 < \infty$ , a sufficient condition for  $n^{-1}ER_{1n} \rightarrow 0$  is to require that the partition satisfy  $\sum_{n=0}^{\infty} n^{\gamma} \pi(\lambda_n) < \infty$  for some  $\gamma > 1/\alpha$ , where  $\pi$  is the ergodic measure of  $X$ .

### 3. Some Extensions

The analysis of Section 2 can be easily extended to cover the case in which  $\{X_n : n \geq 0\}$  is a Harris recurrent Markov chain (see Revuz (1975), p. 75, for the definition). For any initial distribution for  $X_0$ , a probability space can be constructed which supports both the Markov chain  $\{X_n : n \geq 0\}$  and a sequence  $\{T_n : n \geq 1\}$  of random times, and which satisfy:

- i.)  $\{X_{T_n+k} : k \geq 0\}$  has an identical distribution for each  $n \geq 1$
- ii.)  $\mathcal{B}(X_j : j < T_n)$  is independent of  $\mathcal{B}(X_j : j \geq T_{n+1})$  for  $n \geq 1$ ;

see Niemi and Mummelin (1982) for details of the construction. The Markov chain  $X$  can therefore be analyzed as a stochastic process analog of a 1-dependent sequence of identically distributed random variables. To extend the results of Section 2 to our current setting, it is necessary to obtain a moment bound similar to (2.4). Retaining the notation of Section 2, we note that the 1-dependence yields the bound

$$\begin{aligned}
 EW_{2m+1} &= \sum_{n=0}^{\infty} P\{T_{2m+1} < S(n) < T_{2m+2}\} \\
 (3.1) \quad &< \sum_{n=0}^{\infty} \prod_{j=0}^{m-1} P\{Y_{T_{2j}} \neq n, \dots, Y_{T_{2j+1}} \neq n\} \cdot P\{S(n) < T_2\} \\
 &= \sum_{n=0}^{\infty} P\{S(n) > T_2\}^m P\{S(n) < T_2\}.
 \end{aligned}$$

Since Lemma 2.6 clearly continues to hold, systematic application of (3.1) proves that Theorem 2.7 and its Corollary 2.8 remain valid in the Harris chain setting. Further verification, based also on (3.1), proves that Proposition 2.18 also works for Harris chains.

Our second extension concerns the case where the collection  $\Gamma = \{A_n : n \geq 0\}$  is not a partition of  $E$ . For a given family  $\Gamma$  of sets satisfying  $\bigcup_{A \in \Gamma} A = E$ , let

$$(3.2) \quad \phi(\Gamma, \Lambda) = \text{card}\{A \cap \Lambda : A \in \Gamma\}$$

for  $\Lambda \subseteq E$ . If  $\Gamma$  is a countable partition of  $E$  and  $\Lambda_n = \{X_0, \dots, X_n\}$ , then  $\phi(\Gamma, \Lambda_n) = R(0, n) + 1$ , where  $R(0, n)$  is the range process associated with the partition  $\Gamma$ . Thus, (3.2) legitimately generalizes the range sequence studied in Section 2. In the case that  $\Gamma$  is not a partition,  $\phi(\Gamma, \Lambda_n)$  can potentially be of magnitude  $2^n$ , and hence it is not reasonable to expect that  $\phi(\Gamma, \Lambda_n)/n$  will always converge. Instead, it is natural to study  $\log \phi(\Gamma, \Lambda_n)/n$ .

Set  $\phi_{ik} = \phi(\Gamma, \Lambda_{ik})$  for  $0 < i < k$  where  $\Lambda_{ik} = \{X_{T_i}, \dots, X_{T_k-1}\}$ .

(3.3) Lemma. For  $0 < i < j < k$ ,  $\log \phi_{ik} \leq \log \phi_{ij} + \log \phi_{jk}$ .

Proof. The argument follows that used by Steele (1978). Note that  $A \cap \Lambda_{ik} = (A \cap \Lambda_{ij}) \cup (A \cap \Lambda_{jk})$ , so that there are fewer sets of the form of the form  $A \cap \Lambda_{ik}$  than pairs of sets  $A \cap \Lambda_{ij}$ ,  $A \cap \Lambda_{jk}$ . It follows that  $\phi_{ik} \leq \phi_{ij} \phi_{jk}$ . ||

The following theorem is an easy consequence.

(3.4) Theorem. Let  $X$  be a delayed regenerative sequence satisfying  $E \log \phi_{12} < \infty$ . Then,  $(\log \phi_{1n})/n \rightarrow V(\Gamma)$  a.s. as  $n \rightarrow \infty$ , where  $V(\Gamma)$  is a finite constant.

Proof. Since  $\{\log \phi_{ik} : 1 \leq i < k\}$  has the same distribution as  $\{\log \phi_{i+1, k+1} : 1 \leq i < k\}$ , Lemma 3.3 and our moment hypothesis allow one to apply the subadditive ergodic theorem. As a consequence,  $(\log \phi_{in})/n \rightarrow V(\Gamma)$  a.s. as  $n \rightarrow \infty$ , for each  $i \geq 1$ . Since the limit holds for each  $i \geq 1$ , it follows that  $V(\Gamma)$  is a tail random variable, in the sense that  $V(\Gamma)$  is independent of  $\beta(X_j : j < T_i)$  for each  $i \geq 1$ . Hence, a zero-one law applies and  $V(\Gamma)$  is constant a.s. ||

(3.5) Corollary. Let  $X$  be a delayed regenerative sequence satisfying  $E\tau_1 < \infty$ . Then  $(\log \phi(\Gamma, \Lambda_n))/n \rightarrow V(\Gamma)/E\tau_1$  a.s. as  $n \rightarrow \infty$ .

Proof. Since  $\log \phi_{12} \leq \tau_1 \log 2$ , the moment hypothesis of Theorem 3.4 is satisfied, so

$$(3.6) \quad (\log \phi_{1n})/n \rightarrow V(\Gamma) \text{ a.s.}$$

as  $n \rightarrow \infty$ .

We now show that if  $W_1 \subseteq W_2$ , then  $\phi(\Gamma, W_1) \leq \phi(\Gamma, W_2)$ . Observe that two sets  $A_1 \cap W$  are distinct if and only if  $(A_1 \Delta A_2) \cap W \neq \emptyset$ , where  $\Delta$  denotes symmetric difference. Thus, if  $W_1 \subseteq W_2$ , there are more sets of the form  $A \cap W_2$  than  $A \cap W_1$ , proving that  $\phi(\Gamma, W_1) \leq \phi(\Gamma, W_2)$ . Since  $A_{0n} \supseteq A_{1n}$ , it follows that  $\phi_{0n} \geq \phi_{1n}$ . Thus, Lemma 3.3 proves that  $\log \phi_{0n}$  can be squeezed by  $\log \phi_{1n}$ :

$$(3.7) \quad \log \phi_{1n} \leq \log \phi_{0n} \leq \log \phi_{01} + \log \phi_{1n}.$$

Let  $N(m) = \max\{k : T_k \leq m\}$  and observe that

$$(3.8) \quad A_{T_{N(m)}-1} \subseteq A_m \subseteq A_{T_{N(m)}+1}.$$

Relations (3.7) and (3.8) allows us to "squeeze"  $\log \phi(\Gamma, A_m)$  via

$$(3.9) \quad \log \phi_{1, N(m)} \leq \log \phi(\Gamma, A_m) \leq \log \phi_{01} + \log \phi_{1, N(m)+1}.$$

Dividing through in (3.9) by  $N(m)$ , using (3.6), and exploiting the fact that

$N(m)/m \rightarrow 1/E\tau_1$  a.s. yields the desired conclusion.  $\square$

To conclude this section, we wish to point out that the constant  $V(\Gamma)/E\tau_1$  has been extensively studied in a rather different context. Vapnik and Chervonenkis (1971) showed that if  $\{X_n : n \geq 0\}$  is a sequence of independent and identically distributed variates, then the empirical discrepancy function associated with  $\Gamma$ , namely

$$\frac{1}{n} \sup_{A \in \Gamma} \left| \sum_{i=1}^n I(X_i \in A) - nP\{X_1 \in A\} \right|$$

converges to zero a.s. if and only if  $V(\Gamma)/E\tau_1$  vanishes (see also [13]); the constant  $V(\Gamma)/E\tau_1$  is called the Vapnik-Chervonenkis entropy associated with  $\Gamma$ .

#### References

- [1]. Chosid, L. and Isaac, R. (1978). On the range of recurrent Markov chains. Ann. Probability 6, 680-687.
- [2]. Chosid, L. and Isaac, R. (1980). Correction to "On the range of recurrent Markov chains". Ann. Probability 8, 1000.
- [3]. Chung, K. L. (1967). Markov Chains with Stationary Transition Probabilities. Springer-Verlag, New York.
- [4]. Çinlar, E. (1975). Introduction to Stochastic Processes. Prentice-Hall, Englewood Cliffs, NJ.
- [5]. Derriennic, Y. (1980). Quelques applications du Theoreme Ergodique sous-additif in Journées sur les marches aleatoires Nancy 1979. Asterique 74, 183-201.
- [6]. Dvoretzky, A. and Erdos, P. (1951). Some problems on random walk in space. Second Berkeley Symp. Math. Statist. Prob. 353-368.
- [7]. Feller, W. (1971). An Introduction to Probability Theory and its Applications, Vol. II. Wiley, New York.
- [8]. Jain, N. C. and Pruitt, W. E. (1972). The range of random walk. Sixth Berkeley Symp. Math. Statist. Prob. 3, 31-50.
- [9]. Kingman, J. F. C. (1973). Subadditive ergodic theory. Ann. Probability 1 883-909.
- [10]. Niemi, S. and Nummelin, E. (1982). Central limit theorems for Markov random walks. Commentationes Physico-Mathematicae, 54, Societas Scientiarum Fennica, Helsinki.
- [11]. Revuz, D. (1975). Markov Chains. North-Holland, Amsterdam.
- [12]. Spitzer, F. (1976). Principles of Random Walk. Springer, New York.
- [13]. Steele, J. M. (1978). Empirical discrepancies and subadditive processes. Ann. Probability 6, 118-127.
- [14]. Vapnik, V. N. and Chervonenkis, A. Y. (1971). On the uniform convergence of relative frequencies of events to their probabilities. Theory Probability Appl. 16, 264-280.

PWG/jvs

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER #2661	2. GOVT ACCESSION NO. AD-A141652	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle)  On the Range of a Regenerative Sequence		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s)  Peter W. Glynn		8. CONTRACT OR GRANT NUMBER(s)  DAAG29-80-C-0041
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 5 - Optimization and Large Scale Systems
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, North Carolina 27709		12. REPORT DATE March 1984
		13. NUMBER OF PAGES 12
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)  UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  regenerative process, range, subadditive process, Markov chain		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  For a given countable partition of the range of a regenerative sequence $\{X_n : n \geq 0\}$ , let $R_n$ be the number of distinct sets in the partition visited by $X$ up to time $n$ . We study convergence issues associated with the range sequence $\{R_n : n \geq 0\}$ . As an application, we generalize a theorem of Chosid and Isaac to Harris recurrent Markov chains.		

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

**DATA  
FILM**